

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 July 2001 (26.07.2001)

PCT

(10) International Publication Number
WO 01/53489 A1

- (51) International Patent Classification⁷: C12N 15/12, C07K 14/705
- (21) International Application Number: PCT/US01/02113
- (22) International Filing Date: 18 January 2001 (18.01.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/176,692 18 January 2000 (18.01.2000) US
- (71) Applicant: LEXICON GENETICS INCORPORATED
[US/US]; 4000 Research Forest Drive, The Woodlands, TX 77381 (US).
- (72) Inventors: TURNER, C., Alexander, Jr.; 67 Winter Wheat Place, The Woodlands, TX 77381 (US). MATHUR, Brian; 12000 Sawmill Road #2014, The Woodlands, TX 77380 (US). FRIEDRICH, Glenn; Breland & Breland, 2207 Hermann Drive, Houston, TX 77004 (US). ZAMBROWICZ, Brian; 18 Firethorne Place, The Woodlands, TX 77382 (US). SANDS, Arthur, T.; 163 Bristol Bend Circle, The Woodlands, TX 77382 (US).
- (74) Agents: ISHIMOTO, Lance, K. et al.; Lexicon Genetics Incorporated, 4000 Research Forest Drive, The Woodlands, TX 77381 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/53489 A1

(54) Title: HUMAN GABA RECEPTOR PROTEINS AND POLYNUCLEOTIDES ENCODING THE SAME

(57) Abstract: Novel human polynucleotide and polypeptide sequences are disclosed that share sequence similarity with animal gamma-amino butyric acid (GABA) receptor subunits and particularly rho3 subunits thereof. These sequences can be used in therapeutic, diagnostic, and pharmacogenomic applications.

HUMAN GABA RECEPTOR PROTEINS AND POLYNUCLEOTIDES ENCODING THE SAME

The present application claims the benefit of U.S. Provisional Application Number 60/176,692 which was filed on January 18, 2000 and is herein incorporated by reference in its entirety.

1. INTRODUCTION

The present invention relates to the discovery, identification, and characterization of novel human polynucleotides encoding proteins that share sequence similarity with animal gamma-amino butyric acid (GABA) receptor subunits. The invention encompasses the described polynucleotides, host cell expression systems, the encoded proteins, fusion proteins, polypeptides and peptides, antibodies to the encoded proteins and peptides, and genetically engineered animals that either lack or over express the disclosed sequences, antagonists and agonists of the proteins, and other compounds that modulate the expression or activity of the proteins encoded by the disclosed sequences that can be used for diagnosis, drug screening, clinical trial monitoring and the treatment of diseases and disorders.

2. BACKGROUND OF THE INVENTION

Membrane proteins play important roles as, *inter alia*, cell surface markers, receptors, and mediators of signal transduction. GABA receptors bind a potent inhibitory neurotransmitter and this interaction serves as a target for a variety of pharmaceutical agents such as benzodiazepines, barbiturates and alcohol.

3. SUMMARY OF THE INVENTION

The present invention relates to the discovery, identification, and characterization of nucleotides that encode novel human proteins, and the corresponding amino acid sequences of these proteins. The novel human proteins (NHPs) described for

the first time herein share structural similarity with membrane receptors such as, but not limited to human and other mammalian GABA receptors.

The novel human nucleic acid sequences described herein,
5 encode alternative proteins/open reading frames (ORFs) of 467,
392, 180, 420, 345, and 133 amino acids in length (see SEQ ID NOS:
2, 4, 6, 8, 10, and 12 respectively).

The invention also encompasses agonists and antagonists of
the described NHPs, including small molecules, large molecules,
10 mutant NHPs, or portions thereof that compete with native NHP,
peptides, and antibodies, as well as nucleotide sequences that can
be used to inhibit the expression of the described NHPs (e.g.,
antisense and ribozyme molecules, and gene or regulatory sequence
replacement constructs) or to enhance the expression of the
15 described NHP sequences (e.g., expression constructs that place
the described sequence under the control of a strong promoter
system), and transgenic animals that express a NHP transgene, or
"knock-outs" (which can be conditional) that do not express a
functional NHP. A gene trapped murine ES cell line has been
20 produced that knocks-out a murine ortholog of the described NHPs.

Further, the present invention also relates to processes for
identifying compounds that modulate, i.e., act as agonists or
antagonists, of NHP expression and/or NHP activity that utilize
purified preparations of the described NHPs and/or NHP product, or
25 cells expressing the same. Such compounds can be used as
therapeutic agents for the treatment of any of a wide variety of
symptoms associated with biological disorders or imbalances.

4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

30 The Sequence Listing provides the sequences of the described
NHP ORFs that encode the described NHP amino acid sequences.

5. DETAILED DESCRIPTION OF THE INVENTION

The NHPs, described for the first time herein, are novel proteins that are expressed in, *inter alia*, human cell lines, human testis, brain, adrenal gland cells, and gene trapped human
5 cells.

The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described sequences,
10 including the specifically described NHPs, and the NHP products; (b) nucleotides that encode one or more portions of the NHPs that correspond to functional domains, and the polypeptide products specified by such nucleotide sequences, including but not limited to the novel regions of any active domain(s); (c) isolated
15 nucleotides that encode mutant versions, engineered or naturally occurring, of the described NHPs in which all or a part of at least one domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including but not limited to soluble proteins and peptides in which all or a portion
20 of the signal sequence is deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of an NHP, or one of its domains (e.g., a receptor or ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic
25 or diagnostic derivatives of the described polynucleotides such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs comprising a sequence first disclosed in the Sequence Listing.

As discussed above, the present invention includes: (a) the
30 human DNA sequences presented in the Sequence Listing (and vectors comprising the same) and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the

Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989, Current
5 Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3) and encodes a functionally equivalent gene product. Additionally contemplated are any nucleotide sequences that hybridize to the complement of a DNA sequence that encodes and
10 expresses an amino acid sequence presented in the Sequence Listing under moderately stringent conditions, e.g., washing in 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989, *supra*), yet still encodes a functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in
15 other species and mutant NHPs whether naturally occurring or engineered (by site directed mutagenesis, gene shuffling, directed evolution as described in, for example, U.S. Patent No. 5,837,458). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

20 Additionally contemplated are polynucleotides encoding NHP ORFs, or their functional equivalents, encoded by polynucleotide sequences that are about 99, 95, 90, or about 85 percent similar or identical to corresponding regions of the nucleotide sequences of the Sequence Listing (as measured by BLAST sequence comparison
25 analysis using, for example, the GCG sequence analysis package (Madison, Wisconsin) using standard default settings).

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP nucleotide sequences. Such
30 hybridization conditions may be highly stringent or less highly stringent, as described above. In instances where the nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such molecules are generally about 16 to about 100 bases long, or about

20 to about 80, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in
5 conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene
10 expression patterns (particularly using a micro array or high-throughput "chip" format). Additionally, a series of the described NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. An oligonucleotide or polynucleotide
15 sequence first disclosed in at least a portion of one or more of the sequences of SEQ ID NOS: 1-12 can be used as a hybridization probe in conjunction with a solid support matrix/substrate (resins, beads, membranes, plastics, polymers, metal or metallized substrates, crystalline or polycrystalline substrates, etc.). Of
20 particular note are spatially addressable arrays (i.e., gene chips, microtiter plates, etc.) of oligonucleotides and polynucleotides, or corresponding oligopeptides and polypeptides, wherein at least one of the biopolymers present on the spatially addressable array comprises an oligonucleotide or polynucleotide
25 sequence first disclosed in at least one of the sequences of SEQ ID NOS: 1-12, or an amino acid sequence encoded thereby. Methods for attaching biopolymers to, or synthesizing biopolymers on, solid support matrices, and conducting binding studies thereon are disclosed in, *inter alia*, U.S. Patent Nos. 5,700,637, 5,556,752,
30 5,744,305, 4,631,211, 5,445,934, 5,252,743, 4,713,326, 5,424,186, and 4,689,405 the disclosures of which are herein incorporated by reference in their entirety.

Addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-12 can be used to identify and characterize the temporal and tissue specific expression of a gene. These addressable arrays incorporate oligonucleotide sequences of sufficient length to confer the required specificity, yet be within the limitations of the production technology. The length of these probes is within a range of between about 8 to about 2000 nucleotides. Preferably the probes consist of 60 nucleotides and more preferably 25 nucleotides from the sequences first disclosed in SEQ ID NOS:1-12.

For example, a series of the described oligonucleotide sequences, or the complements thereof, can be used in chip format to represent all or a portion of the described sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length can partially overlap each other and/or the sequence may be represented using oligonucleotides that do not overlap. Accordingly, the described polynucleotide sequences shall typically comprise at least about two or three distinct oligonucleotide sequences of at least about 8 nucleotides in length that are each first disclosed in the described Sequence Listing. Such oligonucleotide sequences can begin at any nucleotide present within a sequence in the Sequence Listing and proceed in either a sense (5'-to-3') orientation vis-a-vis the described sequence or in an antisense orientation.

Microarray-based analysis allows the discovery of broad patterns of genetic activity, providing new understanding of gene functions and generating novel and unexpected insight into transcriptional processes and biological mechanisms. The use of addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-12 provides detailed information about transcriptional changes involved in a specific pathway, potentially leading to the

identification of novel components or gene functions that manifest themselves as novel phenotypes.

Probes consisting of sequences first disclosed in SEQ ID NOS:1-12 can also be used in the identification, selection and validation of novel molecular targets for drug discovery. The use of these unique sequences permits the direct confirmation of drug targets and recognition of drug dependent changes in gene expression that are modulated through pathways distinct from the drugs intended target. These unique sequences therefore also have utility in defining and monitoring both drug action and toxicity.

As an example of utility, the sequences first disclosed in SEQ ID NOS:1-12 can be utilized in microarrays or other assay formats, to screen collections of genetic material from patients who have a particular medical condition. These investigations can also be carried out using the sequences first disclosed in SEQ ID NOS:1-12 *in silico* and by comparing previously collected genetic databases and the disclosed sequences using computer software known to those in the art.

Thus the sequences first disclosed in SEQ ID NOS:1-12 can be used to identify mutations associated with a particular disease and also as a diagnostic or prognostic assay.

Although the presently described sequences have been specifically described using nucleotide sequence, it should be appreciated that each of the sequences can uniquely be described using any of a wide variety of additional structural attributes, or combinations thereof. For example, a given sequence can be described by the net composition of the nucleotides present within a given region of the sequence in conjunction with the presence of one or more specific oligonucleotide sequence(s) first disclosed in the SEQ ID NOS: 1-12. Alternatively, a restriction map specifying the relative positions of restriction endonuclease digestion sites, or various palindromic or other specific oligonucleotide sequences can be used to structurally describe a

given sequence. Such restriction maps, which are typically generated by widely available computer programs (e.g., the University of Wisconsin GCG sequence analysis package, SEQUENCHER 3.0, Gene Codes Corp., Ann Arbor, MI, etc.), can optionally be
5 used in conjunction with one or more discrete nucleotide sequence(s) present in the sequence that can be described by the relative position of the sequence relative to one or more additional sequence(s) or one or more restriction sites present in the disclosed sequence.

10 For oligonucleotide probes, highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP gene antisense molecules,
15 useful, for example, in NHP gene regulation (for and/or as antisense primers in amplification reactions of NHP nucleic acid sequences). With respect to NHP gene regulation, such techniques can be used to regulate biological functions. Further, such sequences may be used as part of ribozyme and/or triple helix
20 sequences that are also useful for NHP gene regulation.

Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety which is selected from the group including but not limited to
5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil,
25 hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl)uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine,
30 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil,

5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5 5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

10 In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or 15 analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to 20 each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330). Alternatively, double stranded RNA can be used to disrupt the 25 expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate 30 oligonucleotides can be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass

polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual (and periodic updates thereof), Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y.

Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

Further, a NHP gene homolog can be isolated from nucleic acid obtained from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA prepared from human or non-human cell lines or tissue known or suspected to express an allele of a NHP gene.

The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified
5 fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via the screening of a genomic library.

PCR technology can also be used to isolate full length cDNA
10 sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source (*i.e.*, one known, or suspected, to express a NHP sequence). A reverse transcription (RT) reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the
15 amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a complementary primer. Thus, cDNA sequences upstream of the
20 amplified fragment can be isolated. For a review of cloning strategies that can be used, see *e.g.*, Sambrook et al., 1989, *supra*.

A cDNA encoding a mutant NHP gene can be isolated, for example, by using PCR. In this case, the first cDNA strand may be
25 synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying a mutant NHP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide
30 that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in

the art. By comparing the DNA sequence of the mutant NHP allele to that of a corresponding normal NHP allele, the mutation(s) responsible for the loss or alteration of function of the mutant NHP gene product can be ascertained.

5 Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, obesity, vision disorders, high blood pressure, depression, infertility, etc.), or a cDNA library
10 can be constructed using RNA from a tissue known, or suspected, to express a mutant NHP allele. A normal NHP gene, or any suitable fragment thereof, can then be labeled and used as a probe to identify the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP gene sequences can then be purified
15 and subjected to sequence analysis according to methods well known to those skilled in the art.

 Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant NHP allele in an
20 individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue can be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below. (For screening techniques, see,
25 for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.)

 Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins. In
30 cases where a NHP mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to a NHP are likely to cross-react with a corresponding mutant NHP gene product. Library

clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known in the art.

The invention also encompasses (a) DNA vectors that contain
5 any of the foregoing NHP coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculo virus as described in U.S. Patent
10 No. 5,869,336 herein incorporated by reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d) genetically engineered host cells that express an
15 endogenous NHP gene under the control of an exogenously introduced regulatory element (i.e., gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate
20 expression. Such regulatory elements include but are not limited to the cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 adenovirus, the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the major operator and
25 promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast α -mating factors.

The present invention also encompasses antibodies and anti-
30 idiotypic antibodies (including Fab fragments), antagonists and agonists of the NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP gene (transcription factor inhibitors, antisense and ribozyme molecules, or gene or

regulatory sequence replacement constructs), or promote the expression of a NHP (e.g., expression constructs in which NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

5 The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant NHPs or inappropriately expressed NHPs for the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression
10 systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of NHP in the
15 body. The use of engineered host cells and/or animals may offer an advantage in that such systems allow not only for the identification of compounds that bind to the endogenous receptor for an NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

20 Finally, the NHP products can be used as therapeutics. For example, soluble derivatives such as NHP peptides/domains corresponding to the NHPs, NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies
25 (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream targets in a NHP-mediated pathway) can be used to directly treat diseases or disorders. For instance, the administration of an effective amount of soluble NHP, or a NHP-IgFc fusion protein or an anti-
30 idiotypic antibody (or its Fab) that mimics the NHP could activate or effectively antagonize the endogenous NHP receptor. Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these

genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP fusion protein to the body. Nucleotide constructs encoding functional NHPs, mutant NHPs, as well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating biological disorders.

Various aspects of the invention are described in greater detail in the subsections below.

5.1 THE NHP SEQUENCES

The cDNA sequences and the corresponding deduced amino acid sequences of the described NHPs are presented in the Sequence Listing. The NHP nucleotides were obtained from clustered human gene trapped sequences, ESTs, and cDNA isolated from a human testis cell library. The described sequences share structural similarity with GABA receptor proteins, and particularly rho 3 subunits of the GABA receptor.

5.2 NHPS AND NHP POLYPEPTIDES

NHPs, polypeptides, peptide fragments, mutated, truncated, or deleted forms of the NHPs, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include but are not limited to the generation of antibodies, as reagents in diagnostic assays, for the identification of other cellular gene products related to a NHP, as reagents in assays for screening for compounds that can be as pharmaceutical reagents useful in the therapeutic treatment of mental, biological, or medical disorders and diseases. Given the similarity information and expression data, the described NHPs can be targeted (by drugs, oligos, antibodies, etc,) in order to treat disease, or to therapeutically augment the efficacy of therapeutic agents.

The Sequence Listing discloses the amino acid sequences encoded by the described NHP sequences. The NHPs typically display initiator methionines in DNA sequence contexts consistent with a translation initiation site, and a signal sequence
5 characteristic of membrane or secreted proteins.

The NHP amino acid sequences of the invention include the amino acid sequences presented in the Sequence Listing as well as analogues and derivatives thereof. Further, corresponding NHP homologues from other species are encompassed by the invention.
10 In fact, any NHP protein encoded by the NHP nucleotide sequences described above are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is well known, and,
15 accordingly, each amino acid presented in the Sequence Listing, is generically representative of the well known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together
20 with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al. eds., Scientific American Books, New York, NY, herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can encode such amino
25 acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences as judged by any of a number of criteria, including, but not limited to, the ability to bind and cleave a
30 substrate of a NHP, or the ability to effect an identical or complementary downstream pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, transport, etc.). Such functionally equivalent

NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described above, but which result in a silent change, thus producing a functionally equivalent gene product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where, as in the present instance, the NHP peptide or polypeptide is thought to be membrane protein, the hydrophobic regions of the protein can be excised and the resulting soluble peptide or polypeptide can be recovered from the culture media. Such expression systems also encompass engineered host cells that express a NHP, or functional equivalent, *in situ*. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well known to those skilled in the art. However, such engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of the NHP, but to assess biological activity, e.g., in drug screening assays.

The expression systems that can be used for purposes of the invention include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors

- containing NHP nucleotide sequences; yeast (*e.g.*, *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing NHP nucleotide sequences; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus)
- 5 containing NHP sequences; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing NHP nucleotide sequences; or mammalian cell systems
- 10 (*e.g.*, COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter).
- 15 In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large quantity of such a protein is to be produced for the generation of pharmaceutical compositions of or containing NHP, or for raising
- 20 antibodies to a NHP, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which a NHP coding sequence may be ligated
- 25 individually into the vector in frame with the *lacZ* coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Pharmacia or American Type Culture Collection) can also
- 30 be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by

adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

5 In an insect system, *Autographa californica* nuclear polyhydrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. A NHP coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed
10 under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of NHP coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These
15 recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted sequence is expressed (e.g., see Smith et al., 1983, J. Virol. 46:584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression
20 systems may be utilized. In cases where an adenovirus is used as an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome
25 by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (e.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific
30 initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an

entire NHP gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is
5 inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and
10 initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (See Bittner et al., 1987, Methods in Enzymol. 153:516-544).

15 In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of
20 the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To
25 this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in
30 particular, human cell lines.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the NHP sequences described above can be

engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the NHP product.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk⁻, hgp^rt⁻ or ap^rt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hyg^r, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147).

Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺-nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

5.3 ANTIBODIES TO NHP PRODUCTS

Antibodies that specifically recognize one or more epitopes of a NHP, or epitopes of conserved variants of a NHP, or peptide fragments of a NHP are also encompassed by the invention. Such antibodies include but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

The antibodies of the invention may be used, for example, in the detection of NHP in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of NHP. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes for the evaluation of the effect of test compounds on expression and/or activity of a NHP gene product. Additionally, such antibodies can be used in conjunction gene therapy to, for example, evaluate the normal and/or engineered NHP-expressing cells prior to their introduction into

the patient. Such antibodies may additionally be used as a method for the inhibition of abnormal NHP activity. Thus, such antibodies may, therefore, be utilized as part of treatment methods.

- 5 For the production of antibodies, various host animals may be immunized by injection with the NHP, an NHP peptide (e.g., one corresponding to a functional domain of an NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted), functional equivalents of the NHP or mutated variant of the NHP.
- 10 Such host animals may include but are not limited to pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's adjuvant (complete and incomplete), mineral salts such as
- 15 aluminum hydroxide or aluminum phosphate, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by
- 20 combination and or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diptheria toxoid, ovalbumin, cholera toxin or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.
- 25 Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975,
- 30 Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal

Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated *in vitro* or
5 *in vivo*. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, *Proc. Natl. Acad. Sci.*, 81:6851-6855; Neuberger *et al.*, 1984, *Nature*, 312:604-608;
10 Takeda *et al.*, 1985, *Nature*, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different
15 animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patents Nos. 6,075,181 and 5,877,397 and their respective disclosures which are herein incorporated by reference in their entirety. Also favored is the
20 production of fully humanized monoclonal antibodies as described in US Patent No. 6,150,584 and respective disclosures which are herein incorporated by reference in their entirety.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988,
25 *Science* 242:423-426; Huston *et al.*, 1988, *Proc. Natl. Acad. Sci. USA* 85:5879-5883; and Ward *et al.*, 1989, *Nature* 334:544-546) can be adapted to produce single chain antibodies against NHP gene products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid
30 bridge, resulting in a single chain polypeptide.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: the F(ab')₂ fragments which can

be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 5 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using techniques well known to those skilled in the art. (See, e.g., 10 Greenspan & Bona, 1993, FASEB J 7(5):437-444; and Nissinoff, 1991, J. Immunol. 147(8):2429-2438). For example antibodies which bind to a NHP domain and competitively inhibit the binding of NHP to its cognate receptor can be used to generate anti-idiotypes that "mimic" the NHP and, therefore, bind and activate or neutralize a 15 receptor. Such anti-idiotypic antibodies or Fab fragments of such anti-idiotypes can be used in therapeutic regimens involving a NHP mediated pathway.

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as 20 single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing 25 description. Such modifications are intended to fall within the scope of the appended claims. All cited publications, patents, and patent applications are herein incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising at least 24 contiguous bases of nucleotide sequence first disclosed
5 in SEQ ID NO: 1.
2. An isolated nucleic acid molecule comprising a nucleotide sequence that:
 - (a) encodes the amino acid sequence shown in SEQ ID
10 NO: 2; and
 - (b) hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO: 1 or the complement thereof.
- 15 3. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes the amino acid sequence shown in SEQ ID NO: 2.
- 20 4. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes the amino acid sequence shown in SEQ ID NO: 8.

SEQUENCE LISTING

<110> LEXICON GENETICS INCORPORATED

<120> Novel Human GABA Receptor Proteins and
Polynucleotides Encoding the Same

<130> LEX-0123-PCT

<150> US 60/176,692

<151> 2000-01-18

<160> 12

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1404

<212> DNA

<213> Homo sapiens

<400> 1

atggtcctgg	ctttccagtt	agtctccttc	acctacatct	ggatcatatt	gaaaccaa	60
gtttgtgctg	cttctaaca	caagatgaca	caccagcgg	gctcctcttc	aatgaaaca	120
acctgcaaac	aagaaactag	aatgaagaaa	gatgacagta	ccaaagcgcg	gcctcagaaa	180
tatgagcaac	ttctccatat	agaggacaac	gatttcgcaa	tgagacctgg	atttgagggg	240
tctccagtgc	cagtaggtat	agatgtccat	gttgaaagca	ttgacagcat	ttcagagact	300
aacatggact	ttacaatgac	tttttatctc	aggcattact	ggaaagacga	gaggctctcc	360
tttcctagca	cagcaaacaa	aagcatgaca	tttgatcata	gattgaccag	aaagatctgg	420
gtgcctgata	tcttttttgt	ccactctaaa	agatccttca	tccatgatac	aactatggag	480
aatatcatgc	tgcgcgtaca	ccctgatgga	aacgtcctcc	taagtctcag	gataacgggt	540
tcggccatgt	gctttatgga	tttcagcagg	tttcctcttg	acactcaaaa	ttgttctctt	600
gaactggaaa	gctatgccta	caatgaggat	gacctaatgc	tatactggaa	acacggaaac	660
aagtccttaa	atactgaaga	acatatgtcc	ctttctcagt	tcttcattga	agacttcagt	720
gcatttagtg	gattagcttt	ctatagcagc	acaggttgg	acaataggct	tttcatcaac	780
tttggtctaa	ggaggcatgt	tttcttcttt	gtgctgcaaa	cctatttccc	agccatattg	840
atggtgatgc	tttcatgggt	ttcattttgg	attgaccgaa	gagctgttcc	tgcaagagtt	900
tccctgggaa	tcaccacagt	gctgaccatg	tccacaatca	tcactgctgt	gagcgcctcc	960
atgccccagg	tgtctacct	caaggctgtg	gatgtgtacc	tgtgggtcag	ctccctcttt	1020
gtgttcctgt	cagtcattga	gtatgcagct	gtgaactacc	tcaccacagt	ggaagagcgg	1080
aaacaattca	agaagacagg	aaagatttct	aggatgtaca	atattgatgc	agttcaagct	1140
atggcctttg	atggttggtt	ccatgacagc	gagattgaca	tggaccagac	ttccctctct	1200
ctaaactcag	aagacttcat	gagaagaaaa	tcgatatgca	gccccagcac	cgattcatct	1260
cggataaaga	gaagaaaatc	cctaggagga	catgttggtt	gaatcattct	ggaaaacaac	1320
catgtcattg	acacctattc	taggatttta	ttccccattg	tgtatatattt	atttaatttg	1380
ttttactggg	gtgtatatgt	atga				1404

<210> 2

<211> 467

<212> PRT

<213> Homo sapiens

<400> 2

Met Val Leu Ala Phe Gln Leu Val Ser Phe Thr Tyr Ile Trp Ile Ile

1	5	10	15
Leu Lys Pro Asn Val Cys Ala Ala Ser Asn Ile Lys Met Thr His Gln			
20	25	30	
Arg Cys Ser Ser Ser Met Lys Gln Thr Cys Lys Gln Glu Thr Arg Met			
35	40	45	
Lys Lys Asp Asp Ser Thr Lys Ala Arg Pro Gln Lys Tyr Glu Gln Leu			
50	55	60	
Leu His Ile Glu Asp Asn Asp Phe Ala Met Arg Pro Gly Phe Gly Gly			
65	70	75	80
Ser Pro Val Pro Val Gly Ile Asp Val His Val Glu Ser Ile Asp Ser			
85	90	95	
Ile Ser Glu Thr Asn Met Asp Phe Thr Met Thr Phe Tyr Leu Arg His			
100	105	110	
Tyr Trp Lys Asp Glu Arg Leu Ser Phe Pro Ser Thr Ala Asn Lys Ser			
115	120	125	
Met Thr Phe Asp His Arg Leu Thr Arg Lys Ile Trp Val Pro Asp Ile			
130	135	140	
Phe Phe Val His Ser Lys Arg Ser Phe Ile His Asp Thr Thr Met Glu			
145	150	155	160
Asn Ile Met Leu Arg Val His Pro Asp Gly Asn Val Leu Leu Ser Leu			
165	170	175	
Arg Ile Thr Val Ser Ala Met Cys Phe Met Asp Phe Ser Arg Phe Pro			
180	185	190	
Leu Asp Thr Gln Asn Cys Ser Leu Glu Leu Glu Ser Tyr Ala Tyr Asn			
195	200	205	
Glu Asp Asp Leu Met Leu Tyr Trp Lys His Gly Asn Lys Ser Leu Asn			
210	215	220	
Thr Glu Glu His Met Ser Leu Ser Gln Phe Phe Ile Glu Asp Phe Ser			
225	230	235	240
Ala Ser Ser Gly Leu Ala Phe Tyr Ser Ser Thr Gly Trp Tyr Asn Arg			
245	250	255	
Leu Phe Ile Asn Phe Val Leu Arg Arg His Val Phe Phe Phe Val Leu			
260	265	270	
Gln Thr Tyr Phe Pro Ala Ile Leu Met Val Met Leu Ser Trp Val Ser			
275	280	285	
Phe Trp Ile Asp Arg Arg Ala Val Pro Ala Arg Val Ser Leu Gly Ile			
290	295	300	
Thr Thr Val Leu Thr Met Ser Thr Ile Ile Thr Ala Val Ser Ala Ser			
305	310	315	320
Met Pro Gln Val Ser Tyr Leu Lys Ala Val Asp Val Tyr Leu Trp Val			
325	330	335	
Ser Ser Leu Phe Val Phe Leu Ser Val Ile Glu Tyr Ala Ala Val Asn			
340	345	350	
Tyr Leu Thr Thr Val Glu Glu Arg Lys Gln Phe Lys Lys Thr Gly Lys			
355	360	365	
Ile Ser Arg Met Tyr Asn Ile Asp Ala Val Gln Ala Met Ala Phe Asp			
370	375	380	
Gly Cys Tyr His Asp Ser Glu Ile Asp Met Asp Gln Thr Ser Leu Ser			
385	390	395	400
Leu Asn Ser Glu Asp Phe Met Arg Arg Lys Ser Ile Cys Ser Pro Ser			
405	410	415	
Thr Asp Ser Ser Arg Ile Lys Arg Arg Lys Ser Leu Gly Gly His Val			
420	425	430	
Gly Arg Ile Ile Leu Glu Asn Asn His Val Ile Asp Thr Tyr Ser Arg			
435	440	445	
Ile Leu Phe Pro Ile Val Tyr Ile Leu Phe Asn Leu Phe Tyr Trp Gly			

450
Val Tyr Val
465

455

460

<210> 3
<211> 1179
<212> DNA
<213> Homo sapiens

<400> 3
atggtcctgg ctttcagtt agtctccttc acctacatct ggatcatatt gaaaccaaatt 60
gtttgtgctg cttctaacat caagatgaca caccagcggg gctcctcttc aatgaaacaa 120
acctgcaaac aagaaactag aatgaagaaa gatgacagta ccaaagcgcg gcctcagaaa 180
tatgagcaac ttctccatat agaggacaac gatttcgcaa tgagacctgg atttggaggg 240
tctccagtgc cagtaggtat agatgtccat gttgaaagca ttgacagcat ttcagagact 300
aacatggact ttacaatgac tttttatctc aggcattact ggaaagacga gaggctctcc 360
tttcctagca cagcaaacaa aagcatgaca tttgatcata gattgaccag aaagatctgg 420
gtgcctgata tcttttttgt ccactctaaa agatccttca tccatgatac aactatggag 480
aatatcatgc tgcgcgtaca ccctgatgga aacgtcctcc taagtctcag gataacgggt 540
tcggccatgt gctttatgga ttccagcagg ttctctcttg acaactcaaaa ttgttctctt 600
gaactggaaa gctatgccta caatgaggat gacctaatgc tatactggaa acacggaaac 660
aagtccttaa atactgaaga acatatgtcc ctttctcagt tcttcattga agacttcagt 720
gcacttagtg gattagcttt ctatagcagc acaggttggg acaataggct tttcatcaac 780
tttgtgctaa ggaggcatgt tttcttcttt gtgctgcaaa cctatttccc agccatattg 840
atggtgatgc tttcatgggt ttcattttgg attgaccgaa gagctgttcc tgcaagagtt 900
tccctgggaa tcaccacagt gctgaccatg tccacaatca tcaactgctgt gagcgctcc 960
atgccccagg tgctctacct caaggctgtg gatgtgtacc tgtgggtcag ctccctcttt 1020
gtgttctgt cagtcattga gtatgcagct gtgaactacc tcaccacagt ggaagagcgg 1080
aaacaattca agaagacagg aaaggtacag ccttgctctg actatcagat cccttgggga 1140
atgtggaaaa gactaccctt atctattgcc ctctcttga 1179

<210> 4
<211> 392
<212> PRT
<213> Homo sapiens

<400> 4
Met Val Leu Ala Phe Gln Leu Val Ser Phe Thr Tyr Ile Trp Ile Ile
1 5 10 15
Leu Lys Pro Asn Val Cys Ala Ala Ser Asn Ile Lys Met Thr His Gln
20 25 30
Arg Cys Ser Ser Ser Met Lys Gln Thr Cys Lys Gln Glu Thr Arg Met
35 40 45
Lys Lys Asp Asp Ser Thr Lys Ala Arg Pro Gln Lys Tyr Glu Gln Leu
50 55 60
Leu His Ile Glu Asp Asn Asp Phe Ala Met Arg Pro Gly Phe Gly Gly
65 70 75 80
Ser Pro Val Pro Val Gly Ile Asp Val His Val Glu Ser Ile Asp Ser
85 90 95
Ile Ser Glu Thr Asn Met Asp Phe Thr Met Thr Phe Tyr Leu Arg His
100 105 110
Tyr Trp Lys Asp Glu Arg Leu Ser Phe Pro Ser Thr Ala Asn Lys Ser
115 120 125
Met Thr Phe Asp His Arg Leu Thr Arg Lys Ile Trp Val Pro Asp Ile
130 135 140
Phe Phe Val His Ser Lys Arg Ser Phe Ile His Asp Thr Thr Met Glu

```

145          150          155          160
Asn Ile Met Leu Arg Val His Pro Asp Gly Asn Val Leu Leu Ser Leu
          165          170          175
Arg Ile Thr Val Ser Ala Met Cys Phe Met Asp Phe Ser Arg Phe Pro
          180          185          190
Leu Asp Thr Gln Asn Cys Ser Leu Glu Leu Glu Ser Tyr Ala Tyr Asn
          195          200          205
Glu Asp Asp Leu Met Leu Tyr Trp Lys His Gly Asn Lys Ser Leu Asn
          210          215          220
Thr Glu Glu His Met Ser Leu Ser Gln Phe Phe Ile Glu Asp Phe Ser
225          230          235          240
Ala Ser Ser Gly Leu Ala Phe Tyr Ser Ser Thr Gly Trp Tyr Asn Arg
          245          250          255
Leu Phe Ile Asn Phe Val Leu Arg Arg His Val Phe Phe Phe Val Leu
          260          265          270
Gln Thr Tyr Phe Pro Ala Ile Leu Met Val Met Leu Ser Trp Val Ser
          275          280          285
Phe Trp Ile Asp Arg Arg Ala Val Pro Ala Arg Val Ser Leu Gly Ile
          290          295          300
Thr Thr Val Leu Thr Met Ser Thr Ile Ile Thr Ala Val Ser Ala Ser
305          310          315          320
Met Pro Gln Val Ser Tyr Leu Lys Ala Val Asp Val Tyr Leu Trp Val
          325          330          335
Ser Ser Leu Phe Val Phe Leu Ser Val Ile Glu Tyr Ala Ala Val Asn
          340          345          350
Tyr Leu Thr Thr Val Glu Glu Arg Lys Gln Phe Lys Lys Thr Gly Lys
          355          360          365
Val Gln Pro Cys Ser Asp Tyr Gln Ile Pro Trp Gly Met Trp Lys Arg
          370          375          380
Leu Pro Leu Ser Ile Ala Leu Ser
385          390

```

<210> 5

<211> 543

<212> DNA

<213> Homo sapiens

<400> 5

```

atggtcctgg ctttccagtt agtctccttc acctacatct ggatcatatt gaaaccaaatt      60
gtttgtgctg tttctaacat caagatgaca caccagcggg gctcctcttc aatgaaacaa      120
acctgcaaac aagaaactag aatgaagaaa gatgacagta ccaaagcgcg gcctcagaaa      180
tatgagcaac ttctccatat agaggacaac gatttcgcaa tgagacctgg atttggaggg      240
tctccagtgc cagtaggtat agatgtccat gttgaaagca ttgacagcat ttcagagact      300
aacatggact ttacaatgac tttttatctc aggcattact ggaaagacga gaggtctctc      360
tttcctagca cagcaaacaa aagcatgaca tttgatcata gattgaccag aaagatctgg      420
gtgcctgata tcttttttgt ccactctaaa agatccttca tccatgatac aactatggag      480
aatatcatgc tgcgcgtaca cctgatgga aacgtcctcc taagtctcag atgcctacaa      540
tga                                                    543

```

<210> 6

<211> 180

<212> PRT

<213> Homo sapiens

<400> 6

Met Val Leu Ala Phe Gln Leu Val Ser Phe Thr Tyr Ile Trp Ile Ile

```

1           5           10           15
Leu Lys Pro Asn Val Cys Ala Ala Ser Asn Ile Lys Met Thr His Gln
20           25           30
Arg Cys Ser Ser Ser Met Lys Gln Thr Cys Lys Gln Glu Thr Arg Met
35           40           45
Lys Lys Asp Asp Ser Thr Lys Ala Arg Pro Gln Lys Tyr Glu Gln Leu
50           55           60
Leu His Ile Glu Asp Asn Asp Phe Ala Met Arg Pro Gly Phe Gly Gly
65           70           75           80
Ser Pro Val Pro Val Gly Ile Asp Val His Val Glu Ser Ile Asp Ser
85           90           95
Ile Ser Glu Thr Asn Met Asp Phe Thr Met Thr Phe Tyr Leu Arg His
100          105          110
Tyr Trp Lys Asp Glu Arg Leu Ser Phe Pro Ser Thr Ala Asn Lys Ser
115          120          125
Met Thr Phe Asp His Arg Leu Thr Arg Lys Ile Trp Val Pro Asp Ile
130          135          140
Phe Phe Val His Ser Lys Arg Ser Phe Ile His Asp Thr Thr Met Glu
145          150          155          160
Asn Ile Met Leu Arg Val His Pro Asp Gly Asn Val Leu Leu Ser Leu
165          170          175
Arg Cys Leu Gln
180

```

<210> 7
 <211> 1263
 <212> DNA
 <213> Homo sapiens

```

<400> 7
atgaagaaag atgacagtac caaagcgcgg cctcagaaat atgagcaact tctccatata      60
gaggacaacg atttcgcaat gagacctgga tttggagggt ctccagtgcc agtaggtata      120
gatgtccatg ttgaaagcat tgacagcatt tcagagacta acatggactt tacaatgact      180
ttttattctca ggcattactg gaaagacgag aggcctctct ttcctagcac agcaaacaaa      240
agcatgacat ttgatcatag attgaccaga aagatctggg tgcctgatat cttttttgtc      300
cactctaaaa gatccttcat ccatgataca actatggaga atatcatgct gcgcgtacac      360
cctgatggaa acgtcctcct aagtctcagg ataacgggtt cggccatgtg ctttatggat      420
ttcagcagggt ttcctcttga cactcaaaat tgttctcttg aactggaaag ctatgcctac      480
aatgaggatg acctaattgct atactggaaa cacggaaaca agtccttaaa tactgaagaa      540
catatgtccc tttctcagtt cttcattgaa gacttcagtg catctagtgg attagctttc      600
tatagcagca caggttggtg caataggctt ttcattcaact ttgtgctaag gaggcattgt      660
ttcttctttg tgctgcaaac ctatttccca gccatattga tgggtgatgct ttcattgggt      720
tcattttgga ttgaccgaag agctgttcct gcaagagttt ccctgggaat caccacagtg      780
ctgaccatgt ccacaatcat cactgctgtg agcgcctcca tgccccagggt gtcctacctc      840
aaggctgtgg atgtgtacct gtgggtcagc tccctctttg tgttctgtc agtcattgag      900
tatgcagctg tgaactacct caccacagtg gaagagcgga aacaattcaa gaagacagga      960
aagatttcta ggatgtacaa tattgatgca gttcaagcta tggcctttga tggttgttac      1020
catgacagcg agattgacat ggaccagact tccctctctc taaactcaga agacttcatg      1080
agaagaaaaa cgattgacag cccagcacc gattcatctc ggataaagag aagaaaaatcc      1140
ctaggaggac atgttggtag aatcattctg gaaaacaacc atgtcattga cacctattct      1200
aggattttat tccccattgt gtatatttta ttttaattgt tttactgggg tgtatatgta      1260
tga                                                    1263

```

<210> 8
 <211> 420
 <212> PRT

<213> Homo sapiens

<400> 8

```

Met Lys Lys Asp Asp Ser Thr Lys Ala Arg Pro Gln Lys Tyr Glu Gln
 1          5          10          15
Leu Leu His Ile Glu Asp Asn Asp Phe Ala Met Arg Pro Gly Phe Gly
 20          25          30
Gly Ser Pro Val Pro Val Gly Ile Asp Val His Val Glu Ser Ile Asp
 35          40          45
Ser Ile Ser Glu Thr Asn Met Asp Phe Thr Met Thr Phe Tyr Leu Arg
 50          55          60
His Tyr Trp Lys Asp Glu Arg Leu Ser Phe Pro Ser Thr Ala Asn Lys
 65          70          75          80
Ser Met Thr Phe Asp His Arg Leu Thr Arg Lys Ile Trp Val Pro Asp
 85          90          95
Ile Phe Phe Val His Ser Lys Arg Ser Phe Ile His Asp Thr Thr Met
 100          105          110
Glu Asn Ile Met Leu Arg Val His Pro Asp Gly Asn Val Leu Leu Ser
 115          120          125
Leu Arg Ile Thr Val Ser Ala Met Cys Phe Met Asp Phe Ser Arg Phe
 130          135          140
Pro Leu Asp Thr Gln Asn Cys Ser Leu Glu Leu Glu Ser Tyr Ala Tyr
 145          150          155          160
Asn Glu Asp Asp Leu Met Leu Tyr Trp Lys His Gly Asn Lys Ser Leu
 165          170          175
Asn Thr Glu Glu His Met Ser Leu Ser Gln Phe Phe Ile Glu Asp Phe
 180          185          190
Ser Ala Ser Ser Gly Leu Ala Phe Tyr Ser Ser Thr Gly Trp Tyr Asn
 195          200          205
Arg Leu Phe Ile Asn Phe Val Leu Arg Arg His Val Phe Phe Phe Val
 210          215          220
Leu Gln Thr Tyr Phe Pro Ala Ile Leu Met Val Met Leu Ser Trp Val
 225          230          235          240
Ser Phe Trp Ile Asp Arg Arg Ala Val Pro Ala Arg Val Ser Leu Gly
 245          250          255
Ile Thr Thr Val Leu Thr Met Ser Thr Ile Ile Thr Ala Val Ser Ala
 260          265          270
Ser Met Pro Gln Val Ser Tyr Leu Lys Ala Val Asp Val Tyr Leu Trp
 275          280          285
Val Ser Ser Leu Phe Val Phe Leu Ser Val Ile Glu Tyr Ala Ala Val
 290          295          300
Asn Tyr Leu Thr Thr Val Glu Glu Arg Lys Gln Phe Lys Lys Thr Gly
 305          310          315          320
Lys Ile Ser Arg Met Tyr Asn Ile Asp Ala Val Gln Ala Met Ala Phe
 325          330          335
Asp Gly Cys Tyr His Asp Ser Glu Ile Asp Met Asp Gln Thr Ser Leu
 340          345          350
Ser Leu Asn Ser Glu Asp Phe Met Arg Arg Lys Ser Ile Cys Ser Pro
 355          360          365
Ser Thr Asp Ser Ser Arg Ile Lys Arg Arg Lys Ser Leu Gly Gly His
 370          375          380
Val Gly Arg Ile Ile Leu Glu Asn Asn His Val Ile Asp Thr Tyr Ser
 385          390          395          400
Arg Ile Leu Phe Pro Ile Val Tyr Ile Leu Phe Asn Leu Phe Tyr Trp
 405          410          415
Gly Val Tyr Val

```

420

<210> 9
 <211> 1038
 <212> DNA
 <213> Homo sapiens

<400> 9
 atgaagaaag atgacagtac caaagcgagg cctcagaaat atgagcaact tctccatata 60
 gaggacaacg atttcgcaat gagacctgga tttggagggt ctccagtgcc agtaggtata 120
 gatgtccatg ttgaaagcat tgacagcatt tcagagacta acatggactt tacaatgact 180
 ttttatctca ggcattactg gaaagacgag aggctctcct ttcctagcac agcaaacaaa 240
 agcatgacat ttgatcatag attgaccaga aagatctggg tgcctgatat cttttttgtc 300
 cactctaaaa gatccttcat ccatgataca actatggaga atatcatgct gcgcgtacac 360
 cctgatggaa acgtcctcct aagtctcagg ataacggttt cggccatgtg ctttatggat 420
 ttcagcaggt ttctcttga cactcaaaat tgttctcttg aactggaaag ctatgcctac 480
 aatgaggatg acctaagtct atactggaaa cacggaaaca agtccttaaa tactgaagaa 540
 catatgtccc tttctcagtt cttcattgaa gacttcagtg catctagtgg attagctttc 600
 tatagcagca caggttggtg caataggctt ttcactcaact ttgtgctaag gaggcagttt 660
 ttcttctttg tgctgcaaac ctatttccca gccatattga tggatgatgct ttcattgggtt 720
 tcatttttga ttgaccgaag agctgttctt gcaagagtgt ccctgggaat caccacagtg 780
 ctgaccatgt ccacaatcat cactgctgtg agcgcctcca tgccccaggt gtcctacctc 840
 aaggctgtgg atgtgtacct gtgggtcagc tccctctttg tgttctgtgc agtcattgag 900
 tatgcagctg tgaactacct caccacagtg gaagagcgga aacaattcaa gaagacagga 960
 aaggtagcgc cttgctctga ctatcagatc ccttggggaa tgtggaaaag actaccctta 1020
 tctattgccc tctcttga 1038

<210> 10
 <211> 345
 <212> PRT
 <213> Homo sapiens

<400> 10
 Met Lys Lys Asp Asp Ser Thr Lys Ala Arg Pro Gln Lys Tyr Glu Gln
 1 5 10 15
 Leu Leu His Ile Glu Asp Asn Asp Phe Ala Met Arg Pro Gly Phe Gly
 20 25 30
 Gly Ser Pro Val Pro Val Gly Ile Asp Val His Val Glu Ser Ile Asp
 35 40 45
 Ser Ile Ser Glu Thr Asn Met Asp Phe Thr Met Thr Phe Tyr Leu Arg
 50 55 60
 His Tyr Trp Lys Asp Glu Arg Leu Ser Phe Pro Ser Thr Ala Asn Lys
 65 70 75 80
 Ser Met Thr Phe Asp His Arg Leu Thr Arg Lys Ile Trp Val Pro Asp
 85 90 95
 Ile Phe Phe Val His Ser Lys Arg Ser Phe Ile His Asp Thr Thr Met
 100 105 110
 Glu Asn Ile Met Leu Arg Val His Pro Asp Gly Asn Val Leu Leu Ser
 115 120 125
 Leu Arg Ile Thr Val Ser Ala Met Cys Phe Met Asp Phe Ser Arg Phe
 130 135 140
 Pro Leu Asp Thr Gln Asn Cys Ser Leu Glu Leu Glu Ser Tyr Ala Tyr
 145 150 155 160
 Asn Glu Asp Asp Leu Met Leu Tyr Trp Lys His Gly Asn Lys Ser Leu
 165 170 175
 Asn Thr Glu Glu His Met Ser Leu Ser Gln Phe Phe Ile Glu Asp Phe

										180								185								190		
Ser	Ala	Ser	Ser	Gly	Leu	Ala	Phe	Tyr	Ser	Ser	Thr	Gly	Trp	Tyr	Asn													
										195			200			205												
Arg	Leu	Phe	Ile	Asn	Phe	Val	Leu	Arg	Arg	His	Val	Phe	Phe	Phe	Val													
										210			215			220												
Leu	Gln	Thr	Tyr	Phe	Pro	Ala	Ile	Leu	Met	Val	Met	Leu	Ser	Trp	Val													
										225			230			235												
Ser	Phe	Trp	Ile	Asp	Arg	Arg	Ala	Val	Pro	Ala	Arg	Val	Ser	Leu	Gly													
										245			250			255												
Ile	Thr	Thr	Val	Leu	Thr	Met	Ser	Thr	Ile	Ile	Thr	Ala	Val	Ser	Ala													
										260			265			270												
Ser	Met	Pro	Gln	Val	Ser	Tyr	Leu	Lys	Ala	Val	Asp	Val	Tyr	Leu	Trp													
										275			280			285												
Val	Ser	Ser	Leu	Phe	Val	Phe	Leu	Ser	Val	Ile	Glu	Tyr	Ala	Ala	Val													
										290			295			300												
Asn	Tyr	Leu	Thr	Thr	Val	Glu	Glu	Arg	Lys	Gln	Phe	Lys	Lys	Thr	Gly													
										305			310			315												
Lys	Val	Gln	Pro	Cys	Ser	Asp	Tyr	Gln	Ile	Pro	Trp	Gly	Met	Trp	Lys													
										325			330			335												
Arg	Leu	Pro	Leu	Ser	Ile	Ala	Leu	Ser																				
										340			345															

```
<210> 11
<211> 402
<212> DNA
<213> Homo sapiens
```

```

<400> 11
atgaagaaag atgacagtac caaagcgcgg cctcagaaat atgagcaact tctccatata      60
gaggacaacg atttcgcaat gagacctgga tttggagggt ctccagtgcc agtaggtata      120
gatgtccatg ttgaaagcat tgacagcatt tcagagacta acatggactt tacaatgact      180
ttttatctca ggcattactg gaaagacgag aggctctcct ttccatgacac agcaaaacaa      240
agcatgatcat ttgatcatag attgaccaga aagatctggg tgccatgatc ctttttgtc      300
cactctaaaa gatccttcac ccatgataca actatggaga atatcatgct gcgcgtacac      360
cctgatggaa acgtcctcct aagtctcaga tgccataaat ga                      402

```

```
<210> 12
<211> 133
<212> PRT
<213> Homo sapiens
```

<400> 12																
Met	Lys	Lys	Asp	Asp	Ser	Thr	Lys	Ala	Arg	Pro	Gln	Lys	Tyr	Glu	Gln	
1				5					10					15		
Leu	Leu	His	Ile	Glu	Asp	Asn	Asp	Phe	Ala	Met	Arg	Pro	Gly	Phe	Gly	
			20					25					30			
Gly	Ser	Pro	Val	Pro	Val	Gly	Ile	Asp	Val	His	Val	Glu	Ser	Ile	Asp	
		35				40						45				
Ser	Ile	Ser	Glu	Thr	Asn	Met	Asp	Phe	Thr	Met	Thr	Phe	Tyr	Leu	Arg	
	50				55						60					
His	Tyr	Trp	Lys	Asp	Glu	Arg	Leu	Ser	Phe	Pro	Ser	Thr	Ala	Asn	Lys	
65				70					75					80		
Ser	Met	Thr	Phe	Asp	His	Arg	Leu	Thr	Arg	Lys	Ile	Trp	Val	Pro	Asp	
			85						90				95			
Ile	Phe	Phe	Val	His	Ser	Lys	Arg	Ser	Phe	Ile	His	Asp	Thr	Thr	Met	
			100					105					110			

Glu Asn Ile Met Leu Arg Val His Pro Asp Gly Asn Val Leu Leu Ser
115 120 125
Leu Arg Cys Leu Gln
130

INTERNATIONAL SEARCH REPORT

Intern. Application No.
PCT/US 01/02113

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/12 C07K14/705

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ME BAILEY ET AL.,: "Genetic linkages and radiation hybrid mapping of the three human GABA(C) receptor rho subunit genes : GABR1, GABR2 and GABR3" BIOCHIMICA ET BIOPHYSICA ACTA, vol. 1447, no. 2-3, 28 October 1999 (1999-10-28), pages 307-312, XP000999020	1,2
Y	the whole document	3,4
X	T OGURUSU AND R SHINGAI: "Cloning of a putative gamma-aminobutyric acid (GABA) receptor subunit rho 3 cDNA" BIOCHIMICA ET BIOPHYSICA ACTA, vol. 1305, no. 1-2, 7 February 1996 (1996-02-07), pages 15-18, XP000999706	1,2
Y	the whole document	3,4
	--- -/-	

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

16 May 2001

Date of mailing of the international search report

06/06/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx: 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Julia, P

INTERNATIONAL SEARCH REPORT

Inter Application No
PCT/US 01/02113

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	H QIAN ET AL.,: "Molecular and pharmacological properties of GABA-rho subunits from white perch retina" J. NEUROBIOL., vol. 37, no. 2, 1998, pages 305-320, XP000996647	1,2
Y	the whole document	3,4
X	GR. CUTTING ET AL. : "Cloning of the gamma-aminobutyric acid (GABA) RHO-1 cDNA: a GABA receptor subunit highly expressed in the retina" PROC. NATL. ACAD. SCI. USA, vol. 88, April 1991 (1991-04), pages 2673-2677, XP002167521	1,2
Y	the whole document	3,4
X	GR CUTTING ET AL.,: "Identification of a putative gamma-aminobutyric acid (GABA) receptor subunit rho2 cDNA and colocalization of the genes encoding rho2 (GABRR2) and rho1 (GABRR1) to human chromosome 6q14-q21 and mouse chromosome 4" GENOMICS, vol. 12, no. 4, April 1992 (1992-04), pages 801-806, XP000999703	1,2
Y	the whole document	3,4
Y	K. WEGELIUS ET AL., : "Distribution of GABA receptor rho subunit transcripts in the rat brain" EUROPEAN JOURNAL OF NEUROSCIENCE, vol. 10, January 1998 (1998-01), pages 350-357, XP000996648 the whole document	3,4